

ELECTRICITY AND MAGNETISM

Preliminary Examination

Thursday 08/21/2014

09:00 - 12:00 in P-121

Answer a total of **THREE** questions. If you turn in excess solutions, the ones to be graded will be picked at random.

Each answer must be presented **separately** in an answer book or on sheets of paper stapled together. Make sure you clearly indicate who you are and what is the problem you are solving on each book/sheet of paper. Double-check that you include everything you want graded, and nothing else.


You are allowed to use a result stated in one part of a problem in the subsequent parts even if you cannot derive it. On the last page you will find some potentially useful formulas.

Problem 1. A dielectric sphere carries a charge density which depends only on the distance from the center, $\rho(r)$. The sphere undergoes purely radial oscillations, i.e. ρ at any time depends only on the distance r . Show that no radiation can occur.

Problem 2. A plane wave is incident (normal incidence) on a medium with two layers. The first layer has an index of refraction n and is of thickness d . The second layer has an index of refraction n_1 and is of infinite thickness. Show that if $n = n_1^{1/2}$ and $d = \frac{\lambda}{4}$ where λ is the wavelength of the wave, no reflected wave is observed.


Problem 3. Consider a conductor of width a in x direction. The conductivity is due to electrons and the charge carrier density is n . The conductor is placed in magnetic field B in positive z direction. When the current I_y in y direction is flowing through the conductor, the Hall voltage V_H is generated between the edges $x = 0$ and $x = a$. Find the magnitude and the direction of the Hall voltage.

(You can think of charge carriers as moving with constant drift velocity v_d parallel to the direction of the current.)



Problem 4. Two infinite parallel conducting planes, separated by distance a carry uniform current density J in opposite directions.

- a). Determine the magnetic field between the two planes.
- b). What is the pressure of magnetic field on one plane due to the current in the other?



Standard vector operations in three common coordinate systems

Cartesian coordinates x, y, z

$$\begin{aligned}\nabla &= \hat{\mathbf{e}}_x \frac{\partial}{\partial x} + \hat{\mathbf{e}}_y \frac{\partial}{\partial y} + \hat{\mathbf{e}}_z \frac{\partial}{\partial z} \\ \nabla \cdot \mathbf{A} &= \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} \\ \nabla \times \mathbf{A} &= \hat{\mathbf{e}}_x \left(\frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} \right) + \hat{\mathbf{e}}_y \left(\frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} \right) + \hat{\mathbf{e}}_z \left(\frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right) \\ \nabla^2 &= \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}\end{aligned}$$

cylindrical coordinates ρ, ϕ, z

$$\begin{aligned}\nabla &= \hat{\mathbf{e}}_\rho \frac{\partial}{\partial \rho} + \hat{\mathbf{e}}_\phi \frac{1}{\rho} \frac{\partial}{\partial \phi} + \hat{\mathbf{e}}_z \frac{\partial}{\partial z} \\ \nabla \cdot \mathbf{A} &= \frac{1}{\rho} \frac{\partial}{\partial \rho} (\rho A_\rho) + \frac{1}{\rho} \frac{\partial A_\phi}{\partial \phi} + \frac{\partial A_z}{\partial z} \\ \nabla \times \mathbf{A} &= \hat{\mathbf{e}}_\rho \left[\frac{1}{\rho} \frac{\partial A_z}{\partial \phi} - \frac{\partial A_\phi}{\partial z} \right] + \hat{\mathbf{e}}_\phi \left[\frac{\partial A_\rho}{\partial z} - \frac{\partial A_z}{\partial \rho} \right] + \hat{\mathbf{e}}_z \frac{1}{\rho} \left[\frac{\partial}{\partial \rho} (\rho A_\phi) - \frac{\partial A_\rho}{\partial \phi} \right] \\ \nabla^2 &= \frac{1}{\rho} \frac{\partial}{\partial \rho} \rho \frac{\partial}{\partial \rho} + \frac{1}{\rho^2} \frac{\partial^2}{\partial \phi^2} + \frac{\partial^2}{\partial z^2}\end{aligned}$$

spherical polar coordinates r, θ, ϕ

$$\begin{aligned}\nabla &= \hat{\mathbf{e}}_r \frac{\partial}{\partial r} + \hat{\mathbf{e}}_\theta \frac{1}{r} \frac{\partial}{\partial \theta} + \hat{\mathbf{e}}_\phi \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \\ \nabla \cdot \mathbf{A} &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 A_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta A_\theta) + \frac{1}{r \sin \theta} \frac{\partial A_\phi}{\partial \phi} \\ \nabla \times \mathbf{A} &= \hat{\mathbf{e}}_r \frac{1}{r \sin \theta} \left[\frac{\partial}{\partial \theta} (\sin \theta A_\phi) - \frac{\partial A_\theta}{\partial \phi} \right] + \hat{\mathbf{e}}_\theta \left[\frac{1}{r \sin \theta} \frac{\partial A_r}{\partial \phi} - \frac{1}{r} \frac{\partial A_r}{\partial \theta} \right] + \hat{\mathbf{e}}_\phi \frac{1}{r} \left[\frac{\partial}{\partial r} (r A_\theta) - \frac{\partial A_r}{\partial \theta} \right] \\ \nabla^2 &= \frac{1}{r^2} \frac{\partial}{\partial r} r^2 \frac{\partial}{\partial r} + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \sin \theta \frac{\partial}{\partial \theta} + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2}{\partial \phi^2} \\ &\quad \left[\frac{1}{r^2} \frac{\partial}{\partial r} r^2 \frac{\partial}{\partial r} \equiv \frac{1}{r} \frac{\partial^2}{\partial r^2} r \right]\end{aligned}$$