

# THE SURFACE AREA ARE AND THE VOLUME OF N-DIMENSIONAL SPHERE

SPRING SEMESTER 2017

[http://www.phys.uconn.edu/~rozman/Courses/P2400\\_17S/](http://www.phys.uconn.edu/~rozman/Courses/P2400_17S/)



Last modified: May 5, 2017

## 1 Introduction

The volume of n-dimensional sphere of radius  $r$  is proportional to  $r^n$ ,

$$V_n(r) = v(n) r^n, \quad (1)$$

where the proportionality constant,  $v(n)$ , is the volume of the n-dimensional unit sphere.

The surface area of n-dimensional sphere of radius  $r$  is proportional to  $r^{n-1}$ .

$$S_n(r) = s(n) r^{n-1}, \quad (2)$$

where the proportionality constant,  $s(n)$ , is the surface area of the n-dimensional unit sphere.

The n-dimensional sphere is a union of concentric spherical shells:

$$dV_n(r) = S_n(r) dr \quad (3)$$

Therefore the surface area and the volume are related as following:

$$V_n(R) = \int_0^R S_n(r) dr = s(n) \int_0^R r^{n-1} dr = \frac{s(n)}{n} R^n. \quad (4)$$

The surface area and the volume of the unit sphere are related as following:

$$v(n) = \frac{s(n)}{n}. \quad (5)$$

Consider the integral

$$I_n = \int_{-\infty}^{\infty} e^{-x_1^2 - x_2^2 - \dots - x_n^2} dV_n = \int_0^{\infty} e^{-r^2} dV_n(r), \quad (6)$$

where  $dV_n$  is the volume element in cartesian coordinates

$$dV_n = dx_1 dx_2 \dots dx_n \quad (7)$$

and

$$dV_n(r) = s(n) r^{n-1} dr \quad (8)$$

is the volume element in spherical coordinates.

Since the integrand in the first integral in Eq. (6) is a product of identical gaussians of one variable each,

$$I_n = \left( \int_{-\infty}^{\infty} e^{-x^2} dx \right)^n = (\sqrt{\pi})^n = \pi^{\frac{n}{2}}. \quad (9)$$

On the other hand, the second integral in Eq. (6), evaluated in spherical coordinates

$$I_n = \int_0^{\infty} e^{-r^2} s(n) r^{n-1} dr = s(n) \int_0^{\infty} e^{-r^2} r^{n-1} dr = \frac{s(n)}{2} \int_0^{\infty} e^{-r^2} (r^2)^{\frac{n}{2}-1} dr^2 \quad (10)$$

$$= \frac{s(n)}{2} \int_0^{\infty} e^{-t} t^{\frac{n}{2}-1} dt = \frac{s(n)}{2} \Gamma\left(\frac{n}{2}\right) \quad (11)$$

Comparing Eq. (9) and Eq. (11), we obtain:

$$s(n) = \frac{2 \pi^{\frac{n}{2}}}{\Gamma\left(\frac{n}{2}\right)}. \quad (12)$$

$$v(n) = \frac{\pi^{\frac{n}{2}}}{\frac{n}{2} \Gamma\left(\frac{n}{2}\right)} = \frac{\pi^{\frac{n}{2}}}{\Gamma\left(\frac{n}{2} + 1\right)} \quad (13)$$

## 2 Coulomb's law in n-dimension

In three dimensions Coulomb's law takes the form

$$E_3(r) = \frac{1}{4\pi\epsilon_0} \frac{Q}{r}, \quad (14)$$

where  $E_3$  is the magnitude of the electric field,  $\epsilon_0$  is the electric constant, and  $r$  is the distance from the point charge of charge  $Q$ . What is Coulomb's law look like in high dimensions?

We assume the Maxwell equations hold in any dimension. Hence Gauss law still holds:

$$\Phi = \frac{Q}{\epsilon_0}, \quad (15)$$

where  $\Phi$  is the electric flux through a closed surface enclosing any volume,  $Q$  is the total charge enclosed within that volume.

Due to the spherical symmetry of the field created by a point charge,

$$\Phi = E_n(r) S_n(r), \quad (16)$$

where  $E_n(r)$  is the radial component of the electric field,  $S_n(r)$  is the surface area of n-dimensional sphere of radius  $r$ .

$$S_n(r) = s(n) r^{n-1} = \frac{2\pi^{\frac{n}{2}}}{\Gamma\left(\frac{n}{2}\right)} r^{n-1}. \quad (17)$$

Comparing Eq. (15) and Eq. (16),

$$E_n(r) = \frac{Q}{\epsilon_0 S_n(r)} = \frac{\Gamma\left(\frac{n}{2}\right)}{2\pi^{\frac{n}{2}} \epsilon_0} \frac{Q}{r^{n-1}}. \quad (18)$$

## References

- [1] Wikipedia, "Volume of an n-ball — Wikipedia, the free encyclopedia," 2016. Accessed: November 25, 2016.