

# CAUCHY'S INTEGRAL THEOREM: EXAMPLES

SPRING SEMESTER 2024

[https://www.phys.uconn.edu/~rozman/Courses/P2400\\_24S/](https://www.phys.uconn.edu/~rozman/Courses/P2400_24S/)

Last modified: February 11, 2024

Cauchy's theorem states that if  $f(z)$  is analytic at all points on and inside a closed complex contour  $C$ , then the integral of the function around that contour vanishes:

$$\oint_C f(z) dz = 0. \quad (1)$$

## 1 A trigonometric integral

**Problem:** Show that

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos(\alpha \phi) [\cos \phi]^{\alpha-1} d\phi = 2^\alpha B(\alpha, \alpha) = 2^\alpha \frac{\Gamma(\alpha)^2}{\Gamma(2\alpha)}. \quad (2)$$

**Solution:**

Recall the definition of Beta function,

$$B(\alpha, \beta) = \int_0^1 x^{\alpha-1} (1-x)^{\beta-1} dx. \quad (3)$$

Consider the integral:

$$J = \oint_C [z(1-z)]^{\alpha-1} dz, \quad \alpha > 1, \quad (4)$$

where the integration is over closed contour shown in Fig. 1.

Since the integrand in Eq. (4) is analytic inside  $C$ ,

$$J = 0. \quad (5)$$

On the other hand,

$$J = J_I + J_{II}, \quad (6)$$

where  $J_I$  is the integral along the segment of the positive real axis,  $0 \leq x \leq 1$ ;  $J_{II}$  is the integral along the circular arc or radius  $R = \frac{1}{2}$  centered at  $z = \frac{1}{2}$ .

Along the real axis  $z = x$ ,  $dz = dx$ , thus

$$J_I = \int_0^1 x^{\alpha-1} (1-x)^{\alpha-1} dx = B(\alpha, \alpha). \quad (7)$$

Along the semi-circular arc

$$z = \frac{1}{2} + \frac{1}{2} e^{i\theta} = \frac{1}{2} e^{i\frac{\theta}{2}} \left( e^{-i\frac{\theta}{2}} + e^{i\frac{\theta}{2}} \right) = e^{i\frac{\theta}{2}} \cos \frac{\theta}{2}, \quad (8)$$


where  $0 < \theta < \pi$ , and

$$1 - z = \frac{1}{2} - \frac{1}{2} e^{i\theta} = -i e^{i\frac{\theta}{2}} \sin \frac{\theta}{2}. \quad (9)$$

Hence,

$$z(1-z) = -i e^{i\theta} \cos \frac{\theta}{2} \sin \frac{\theta}{2} = -\frac{i}{2} e^{i\theta} \sin \theta. \quad (10)$$

Figure 1: Integration contour for Problem 1



$$dz = \frac{i}{2} e^{i\theta} d\theta. \quad (11)$$

Therefore,

$$J_{\text{II}} = -\left(\frac{-i}{2}\right)^\alpha \int_0^\pi e^{i\alpha\theta} (\sin \theta)^{\alpha-1} d\theta = -2^{-\alpha} e^{-i\frac{\pi}{2}\alpha} \int_0^\pi e^{i\alpha\theta} (\sin \theta)^{\alpha-1} d\theta, \quad (12)$$

where we used that

$$-i = e^{-i\frac{\pi}{2}}. \quad (13)$$

Combining Eqs. (5), (6), (7), and (12) we obtain:

$$2^{-\alpha} e^{-i\frac{\pi}{2}\alpha} \int_0^\pi e^{i\alpha\theta} (\sin \theta)^{\alpha-1} d\theta = B(\alpha, \alpha). \quad (14)$$

Transforming the expression on the left as following:

$$\int_0^\pi e^{i\alpha(\theta-\frac{\pi}{2})} (\sin \theta)^{\alpha-1} d\theta = \int_0^\pi e^{i\alpha(\theta-\frac{\pi}{2})} \left[\cos\left(\theta - \frac{\pi}{2}\right)\right]^{\alpha-1} d\theta \quad (15)$$

$$= \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{i\alpha\phi} [\cos \phi]^{\alpha-1} d\phi \quad (16)$$

$$= \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos(\alpha\phi) [\cos \phi]^{\alpha-1} d\phi \quad (17)$$

finally obtain the relation:

$$\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos(\alpha\phi) [\cos \phi]^{\alpha-1} d\phi = 2^\alpha B(\alpha, \alpha) = 2^\alpha \frac{\Gamma(\alpha)^2}{\Gamma(2\alpha)}. \quad (18)$$

## 2 Euler's log-sine integral.

**Problem:** Show that

$$\int_0^{\pi} \log(\sin x) dx = -\pi \log 2. \quad (19)$$

Integral Eq. (19), is called Euler's log-sine integral. It was first evaluated (by Euler) in 1769.

**Solution:**

We start by integrating the function  $f(z)$ ,

$$f(z) = \log(1 - e^{2iz}), \quad (20)$$

along the rectangular contour  $C$  with the corners at  $0, \pi, \pi + iR, iR$ , indented at the corners when necessary (see Fig. 2), and letting  $R \rightarrow \infty$ .

$$J = \oint_C \log(1 - e^{2iz}) dz. \quad (21)$$

On the one hand, the integrand in Eq. (21) is an analytic function inside  $C$ , therefore

$$J = 0. \quad (22)$$

On the other hand,

$$J = J_I + J_{II} + J_{III} + J_{IV} + J_V + J_{VI}, \quad (23)$$

where the subscripts corresponds to integration contours labeled in Fig. 2.

Consider first the integrals  $J_{II}$  and  $J_{IV}$ . The integrand  $f(z)$  is a periodic function with the period  $\pi$ ,

$$f(z) = f(z + \pi). \quad (24)$$

Indeed,

$$f(z + \pi) = \log(1 - e^{2i(z+\pi)}) = \log(1 - e^{2iz} e^{2\pi i}) = \log(1 - e^{2iz}) = f(z). \quad (25)$$

Therefore in  $J_{II}$  and  $J_{IV}$  we have equal integrands but we are integrating in the opposite directions. Therefore,

$$J_{II} = -J_{IV}, \quad (26)$$




Figure 2: Integration contour for Problem 2

or

$$J_{II} + J_{IV} = 0. \quad (27)$$

Next, observe that  $f(z) \rightarrow 0$  as  $y = \text{Im}(z) \rightarrow +\infty$ :

$$f(z) = \log(1 - e^{2i(x+iy)}) = \log(1 - e^{2ix}e^{-y}) \approx -e^{2ix}e^{-y} \rightarrow 0. \quad (28)$$

Therefore,

$$J_{III} = 0. \quad (29)$$

Next, let's show that

$$J_V \equiv \lim_{r \rightarrow 0} \int_{C_V} \log(1 - e^{2iz}) dz = 0. \quad (30)$$

Indeed,  $z = r e^{i\theta}$ ,  $dz = i r e^{i\theta} d\theta$ ,  $0 \leq \theta \leq \frac{\pi}{2}$ :

$$J_V = i \lim_{r \rightarrow 0} r \int_0^{\frac{\pi}{2}} \log(1 - e^{2ire^{i\theta}}) e^{i\theta} d\theta \approx i \lim_{r \rightarrow 0} r \int_0^{\frac{\pi}{2}} \log(-2ire^{i\theta}) e^{i\theta} d\theta = 0. \quad (31)$$

Similarly we can show that

$$J_{VI} = 0. \quad (32)$$

Combining Eqs (22), (23), (27), (29), (30), and (32), we get that

$$J_I = \int_0^{\pi} \log(1 - e^{2ix}) dx = 0. \quad (33)$$

Rewriting the integrand in Eq. (33) as following,

$$\begin{aligned} \log(1 - e^{2ix}) &= \log[e^{ix}(e^{-ix} - e^{ix})] \\ &= \log\left[2(-i)e^{ix}\left(\frac{e^{ix} - e^{-ix}}{2i}\right)\right] = \log\left(2e^{i(x-\frac{\pi}{2})}\sin x\right) \\ &= \log 2 + i\left(x - \frac{\pi}{2}\right) + \log(\sin x), \end{aligned} \quad (34)$$

we obtain

$$\int_0^{\pi} \left[ \log 2 + i\left(x - \frac{\pi}{2}\right) + \log(\sin x) \right] dx = 0, \quad (35)$$

or

$$\int_0^{\pi} \log(\sin x) dx = -\pi \log 2. \quad (36)$$

### 3 Another Euler integral

**Problem:** evaluate the following integral:

$$I(\alpha) = \int_0^{\infty} \frac{\sin(x)}{x^{\alpha}} dx, \quad 0 < \alpha < 1. \quad (37)$$

**Solution:**

Let's consider the following integral:

$$J(\alpha) = \oint_C \frac{e^{iz}}{z^{\alpha}} dz, \quad (38)$$

where the integration contour  $C$  is sketch in Fig. 3.

The integrand in Eq. (38) is an analytic function inside  $C$ , therefore

$$J(\alpha) = 0. \quad (39)$$

On the other hand,

$$J(\alpha) = J_I + J_{II} + J_{III} + J_{IV}. \quad (40)$$

where the subscripts corresponds to integration contours labeled in Fig. 3.

Let's consider  $J_I$ ,  $J_{II}$ ,  $J_{III}$ , and  $J_{IV}$  separately:

$J_I$ : the integration is along the real axis, so  $z = x$ ,  $dz = dx$ ,  $r \leq x \leq R$ :

$$J_I = \lim_{r \rightarrow 0} \lim_{R \rightarrow \infty} \int_r^R \frac{e^{ix}}{x^{\alpha}} dx = \int_0^{\infty} \frac{e^{ix}}{x^{\alpha}} dx, \quad (41)$$


so

$$I(\alpha) = \text{Im } J_I. \quad (42)$$

$J_{\text{II}}$ : the integration is counterclockwise along the quarter-circle of radius  $R$ ,  $z = R e^{i\theta}$ ,  $dz = i R e^{i\theta} d\theta$ ,  $0 \leq \theta \leq \frac{\pi}{2}$ :

$$J_{\text{II}} = \lim_{R \rightarrow \infty} i R^{(1-\alpha)} \int_0^{\frac{\pi}{2}} e^{i R \cos \theta} e^{-R \sin \theta} e^{i(1-\alpha)\theta} d\theta. \quad (43)$$

Figure 3: Integration contour for Problem 3





For the absolute value of  $J_{\text{II}}$  we have the following estimates:

$$|J_{\text{II}}| = \lim_{R \rightarrow \infty} \left| R^{(1-\alpha)} \int_0^{\frac{\pi}{2}} e^{iR \cos \theta} e^{-R \sin \theta} e^{i(1-\alpha)\theta} d\theta \right| \quad (44)$$

$$\leq \lim_{R \rightarrow \infty} R^{(1-\alpha)} \int_0^{\frac{\pi}{2}} |e^{iR \cos \theta} e^{-R \sin \theta} e^{i(1-\alpha)\theta}| d\theta \quad (45)$$

$$= \lim_{R \rightarrow \infty} R^{(1-\alpha)} \int_0^{\frac{\pi}{2}} e^{-R \sin(\theta)} d\theta \leq \lim_{R \rightarrow \infty} R^{(1-\alpha)} \int_0^{\frac{\pi}{2}} e^{-\frac{2R}{\pi}\theta} d\theta \quad (46)$$

$$= \lim_{R \rightarrow \infty} R^{(1-\alpha)} \frac{\pi}{2R} \int_0^R e^{-u} du = \frac{\pi}{2} \lim_{R \rightarrow \infty} R^{-\alpha} (1 - e^{-R}) = 0, \quad (47)$$

where we used the inequalities

$$\sin(\theta) \geq \frac{2}{\pi}\theta \quad \longrightarrow \quad e^{-\sin(\theta)} \leq e^{-\frac{2}{\pi}\theta} \quad \longrightarrow \quad e^{-R \sin(\theta)} \leq e^{-\frac{2R}{\pi}\theta}, \quad (48)$$

that are valid within the integration range  $0 \leq \theta \leq \frac{\pi}{2}$ , and introduce a new integration variable  $u = \frac{2R}{\pi}\theta$ .

Thus,

$$J_{\text{II}} = 0. \quad (49)$$

$J_{\text{III}}$ : the integration is along the imaginary axis, so  $z = iy$ ,  $dz = i dy$ ,  $r \leq y \leq R$ :

$$J_{\text{I}} = \lim_{r \rightarrow 0} \lim_{R \rightarrow \infty} i^{(1-\alpha)} \int_R^r \frac{e^{-y}}{y^\alpha} dy = -e^{i\frac{\pi}{2}(1-\alpha)} \int_0^\infty e^{-y} y^{-\alpha} dy = -e^{i\frac{\pi}{2}(1-\alpha)} \Gamma(1-\alpha). \quad (50)$$

$J_{\text{IV}}$ : the integration is clockwise along the quarter-circle of radius  $r$ ,  $z = r e^{i\theta}$ ,  $dz = i r e^{i\theta} d\theta$ ,  $0 \leq \theta \leq \frac{\pi}{2}$ :

$$J_{\text{IV}} = \lim_{r \rightarrow 0} i r^{(1-\alpha)} \int_{\frac{\pi}{2}}^0 e^{i r e^{i\theta}} e^{i(1-\alpha)\theta} d\theta \approx -\lim_{r \rightarrow 0} i r^{(1-\alpha)} \int_0^{\frac{\pi}{2}} e^{i(1-\alpha)\theta} d\theta = 0. \quad (51)$$

Combining Eqs. (39), (40), and (52), we get

$$J_I = e^{i\frac{\pi}{2}(1-\alpha)} \Gamma(1-\alpha). \quad (52)$$

Taking the imaginary part, and using Eq. (42), we obtain

$$\int_0^\infty \frac{\sin(x)}{x^\alpha} dx = \sin\left(\frac{\pi}{2}(1-\alpha)\right) \Gamma(1-\alpha). \quad (53)$$

For the case  $\alpha = 1$ ,

$$\int_0^\infty \frac{\sin(x)}{x} dx = \lim_{\alpha \rightarrow 1} \sin\left(\frac{\pi}{2}(1-\alpha)\right) \Gamma(1-\alpha) \approx \lim_{\alpha \rightarrow 1} \frac{\pi}{2}(1-\alpha) \Gamma(1-\alpha) = \frac{\pi}{2} \lim_{\alpha \rightarrow 1} \Gamma(2-\alpha) = \frac{\pi}{2} \Gamma(1) = \frac{\pi}{2}.$$

## 4 Fresnel integrals

**Problem:** Assuming that the value of the Gaussian integral is known,

$$I = \int_0^\infty e^{-x^2} dx = \frac{\sqrt{\pi}}{2}, \quad (54)$$

evaluate the Fresnel integrals,

$$C = \int_0^\infty \cos(x^2) dx \quad (55)$$

and

$$S = \int_0^\infty \sin(x^2) dx. \quad (56)$$

The integrals  $C$  and  $S$  are named after the Fresnel (French physicist, 1788-1827). They were first evaluated by Euler in 1781.

**Solution:**

Let's pack  $C$  and  $S$  together:

$$F \equiv C + iS = \int_0^\infty [\cos(x^2) + i\sin(x^2)] dx = \int_0^\infty e^{ix^2} dx, \quad (57)$$

such that

$$C = \operatorname{Re} F \quad (58)$$

and

$$S = \operatorname{Im} F. \quad (59)$$

Consider the integral

$$J = \int_C e^{iz^2} dz, \quad (60)$$

where  $C$  is the contour in the complex plane shown in Fig. 4.

Since the integrand in Eq. (61) is analytic inside  $C$ ,

$$J = 0. \quad (61)$$

On the other hand,

$$J = J_I + J_{II} + J_{III}, \quad (62)$$

where  $J_I$  is the integral along the positive real axis,  $J_{II}$  is the integral along the circular arc of radius  $R \rightarrow \infty$ ,  $0 \leq \theta \leq \frac{\pi}{4}$ , and  $J_{III}$  is the integral from infinity to the origin along the ray that makes the angle  $\theta = \frac{\pi}{4}$  with the real axis.

Let's consider  $J_I$ ,  $J_{II}$ , and  $J_{III}$  separately:

$J_I$ : the integration is along the real axis, so  $z = x$ ,  $dz = dx$ ,  $0 \leq x \leq \infty$ :




Figure 4: Integration contour for Problem 4

$$J_I = \int_{C_I} e^{iz^2} dz = \int_0^\infty e^{ix^2} dx = F. \quad (63)$$

$J_{II}$ : the integration is along the circular arc of radius  $R$  so  $z = Re^{i\theta}$ ,  $dz = iRe^{i\theta}d\theta$ ,  $z^2 = R^2 e^{2i\theta} = R^2 (\cos(2\theta) + i\sin(2\theta))$ ,  $0 \leq \theta \leq \frac{\pi}{4}$ :

$$J_{II} = \int_{C_{II}} e^{iz^2} dz = iR \int_0^{\frac{\pi}{4}} e^{iR^2 \cos(2\theta)} e^{-R^2 \sin(2\theta)} d\theta. \quad (64)$$

For the absolute value of  $J_{II}$  we have the following estimates:

$$|J_{II}| = \left| R \int_0^{\frac{\pi}{4}} e^{iR^2 \cos(2\theta)} e^{-R^2 \sin(2\theta)} d\theta \right| \leq R \int_0^{\frac{\pi}{4}} \left| e^{iR^2 \cos(2\theta)} e^{-R^2 \sin(2\theta)} \right| d\theta \quad (65)$$

$$= R \int_0^{\frac{\pi}{4}} e^{-R^2 \sin(2\theta)} d\theta = \frac{R}{2} \int_0^{\frac{\pi}{2}} e^{-R^2 \sin(\phi)} d\phi < \frac{R}{2} \int_0^{\frac{\pi}{2}} e^{-\frac{2R^2}{\pi}\phi} d\phi \quad (66)$$

$$= \frac{R}{2} \frac{\pi}{2R^2} \int_0^{R^2} e^{-u} du = \frac{\pi}{4R} (1 - e^{-R^2}) < \frac{\pi}{4R}, \quad (67)$$

where we introduced a new integration variable  $\phi = 2\theta$ , used the inequalities

$$\sin(\phi) \geq \frac{2}{\pi}\phi \quad \longrightarrow \quad e^{-\sin(\phi)} \leq e^{-\frac{2}{\pi}\phi} \quad \longrightarrow \quad e^{-R^2 \sin(\phi)} \leq e^{-\frac{2R^2}{\pi}\phi}, \quad (68)$$

that are valid within the integration range  $0 \leq \phi \leq \frac{\pi}{2}$ , and introduce a new integration variable  $u = \frac{2R^2}{\pi}\phi$ .

Thus we obtained that

$$|J_{II}| < \frac{\pi}{4R}. \quad (69)$$

Therefore,

$$J_{II} = 0 \quad (70)$$

as  $R \rightarrow \infty$ .

$J_{\text{III}}$ : the integration is along the ray making the angle  $\frac{\pi}{4}$  with the real axis so  $z = r e^{i\frac{\pi}{4}}$ ,  
 $z^2 = r^2 e^{i\frac{\pi}{2}} = i r^2$ ,  $dz = e^{i\frac{\pi}{4}} dr$ ,  $0 \leq r < \infty$ .

$$J_{\text{III}} = \int_{C_{\text{III}}} e^{i z^2} dz = e^{i\frac{\pi}{4}} \int_0^\infty e^{-r^2} dr = -e^{i\frac{\pi}{4}} \int_0^\infty e^{-r^2} dr = -e^{i\frac{\pi}{4}} \frac{\sqrt{\pi}}{2}. \quad (71)$$

Combining Eqs. (62), (64), (71), and (72) we obtain:

$$F = \frac{\sqrt{\pi}}{2} e^{i\frac{\pi}{4}}. \quad (72)$$

Finally, the Fresnel's integrals are:

$$C = \operatorname{Re} F = \frac{\sqrt{\pi}}{2} \cos\left(\frac{\pi}{4}\right) = \sqrt{\frac{\pi}{8}} \quad (73)$$

and

$$S = -\operatorname{Im} F = -\frac{\sqrt{\pi}}{2} \sin\left(\frac{\pi}{4}\right) = -\sqrt{\frac{\pi}{8}}. \quad (74)$$